arXiv:1206.4956v1 [quant-ph] 21 Jun 2012

Large deviations and quantum dynamical phase transitions for the atom maser

Merlijn van Horssen and Mădălin Guţă

University of Nottingham, School of Mathematical Sciences
University Park, NG7 2RD Nottingham, UK

June 2012

Abstract

Large deviations is a framework for rigorously studying the probabilities of rare events in terms of a rate function. We study the atom maser model in the context of large deviations, following the work done by Garrahan, Armour and Lesanovsky [1] which relates this rate function to the occurrence of quantum dynamical phase transitions. We show that a large deviations principle holds for the associated classical counting process. This is done by deforming the Lindblad generator for the cavity dynamics and analysing the spectral properties of the associated deformed quantum dynamical semigroup. In particular, the large deviations rate function is given by the spectral bound of the generator. We use the rate function to discuss metastable behaviour of the stationary distribution of the counting process and its connection to quantum dynamical phase transitions.

1 Introduction

In this paper we study the atom counting statistics associated to the atom maser model. In particular, we are interested in whether any quantum dynamical phase transitions occur, using a large deviations approach. Previous work was done in [1–6]; in this paper, we expand on the mathematical background to the results obtained in [1]. The large deviations approach originated in non-equilibrium statistical mechanics [5], where transitions between 'active' and 'passive' dynamical phases were analysed using a large deviations rate function. This rate function plays the role of a free energy in statistical mechanics, and this analogy is used to find dynamical phase transitions as non-analyticities in the rate function.

In general, the large deviations approach involves quantifying rare events; in this case, the events are the trajectories followed by the state of the atom maser cavity. These trajectories can be characterised in term of the number of jumps between energy levels per unit time, and in this sense trajectories can be called active or passive. Trajectories which are highly active or highly passive are considered rare, and the large deviations approach is used to quantify their occurrence. The phase transitions we consider are between values of the dynamical parameters for which active (or passive) trajectories suddenly become common. These points are closely related to the large deviations rate function, which quantifies the occurrence of such trajectories.

In the atom maser model, atoms are prepared in a known state, and interact with a resonant cavity (which is in contact with a thermal bath) one at a time. We link properties of the stationary state of the cavity to the counting statistics of the output atoms. Our main result is that a large deviations principle holds for the counting process associated to atoms emerging from the cavity in the ground state. This means that quantum dynamical phase transitions do not occur in the sense of non-analytic behaviour of the large deviations rate function. The proof of this is based on analysis of the spectral properties of the generator of the dynamics; in particular, we show that a Krein-Rutman theorem applies to the operators in the associated quantum dynamical semigroup.

We note some other previous work on large deviations in quantum systems. In [7] (see also [8]) a large deviation principle is shown to hold for correlated states on quantum spin chains, a related (but finite-dimensional) system; large deviations for quantum Markov semigroups are studied in [9].

This paper is organised as follows. In Section 2 we introduce the background to our problem, starting with the atom maser, where we discuss the master equation and the counting process. The rest of the section consists of the mathematical background, with one part about the theory of large deviations and the other part introducing some operator analysis.

In Section 3 we state our main result, the counting process associated to the atom maser satisfies a large deviations principle; an outline of the proof is given. The proof of our result is found in Section 4. To show that the large deviations principle holds, we prove that a Krein-Rutman theorem holds for the generator of the atom maser dynamics. We discuss the results in Section 5, where we perform a numerical analysis of the problem.

2 Background

Atom maser

The atom maser consists of a sequence of two-level atoms passing successively through a cavity and interacting resonantly with the electromagnetic field inside the cavity. The two-level atoms enter the cavity in a fixed state, chosen here to be the excited state, and for simplicity we assume that only a single atom passes through the cavity at a time. In addition, the cavity is also coupled to a thermal bath which represents the interaction between the (non-ideal) cavity and the environment. The combined effects of the interactions with the atoms and the environment changes the state of the cavity whose time evolution is described by a *quantum Markov semigroup*, in a certain coarse grained approximation; see Refs. [10] and [11] for a mathematical overview, and [12] for the physical derivation of the master equation. Here we will give an intuitive description of the dynamics starting with a simplified discrete time model, with an emphasis on the statistics of measurements performed on the atoms.

The cavity is described by a one mode continuos variables system with Hilbert space $\mathfrak{h} = l^2(\mathbb{N})$ whose canonical basis vectors $(e_n)_{n\geqslant 0}$ represent pure states of fixed number of photons. Therefore, if $\psi \in \mathfrak{h}$ is a pure state, the *photon number distribution* of the cavity is given by $|\langle e_n, \psi\rangle|^2$. In general, (mixed) states are described by density operators i.e. positive trace-class operators $\rho \in \mathcal{T}(\mathfrak{h})$ which are normalised to have unit trace, and the observables are represented by self-adjoint elements of the algebra of bounded operators $\mathcal{B}(\mathfrak{h})$. Recall that the *annihilation operator* a on $\mathfrak{h}$ is defined by

$$a e_n = \begin{cases} \sqrt{n} e_{n-1} & \text{if } n > 0, \\ 0 & \text{if } n = 0, \end{cases}.$$

Its adjoint is the *creation operator* a^* and $N = a^* a$ is the photon number operator. The atom is modelled on the two-dimensional Hilbert space $\mathbb{C}^2$ with standard orthonormal basis $\{|0\rangle, |1\rangle\}$ consisting of the 'ground' and 'excited' states. We denote by σ^* and σ the corresponding raising and lowering operators (i.e. $\sigma^*|0\rangle = |1\rangle$ etc.) The interaction between an atom and the cavity is described by the Jaynes-Cummings hamiltonian

$$H_{\text{int}} = -g(\sigma a^* + \sigma^* a).$$

where g is the coupling constant which for simplicity is considered to be constant across the cavity. The free hamiltonian is

$$H_{\text{free}} = \omega a^* a + \omega \sigma^* \sigma,$$

where ω is the frequency of the resonant mode; however by passing to the interaction picture the effect of the free evolution can be ignored. Therefore if the interaction lasts for a time t_0, the joint evolution is described by the unitary operator $U := \exp(it_0 H_{\text{int}})$ whose action on a product initial state is

$$U : |k\rangle \otimes |1\rangle \mapsto \cos(\phi\sqrt{k+1})|k\rangle \otimes |1\rangle + \sin(\phi\sqrt{k+1})|k+1\rangle \otimes |0\rangle,$$

where $\phi := t_0 g$ is the *accumulated Rabi frequency*. Conditional on the outcome of a measurement in the standard basis of the atom, the cavity remains in state $|k\rangle$ with probability $\cos^2(\phi\sqrt{k+1})$ or it gains an excitation with probability $\sin^2(\phi\sqrt{k+1})$. If we average over the outcome, we obtain a transfer operator T_* acting on the cavity state space

$$\mathcal{T}_*(\rho) = L_1 \rho L_1^* + L_2 \rho L_2^* \tag{1}$$

where the operators L_1 and L_2 are given by

$$L_1 = \cos(\phi\sqrt{aa^*}), \quad L_2 = a^* \frac{\sin(\phi\sqrt{aa^*})}{\sqrt{aa^*}}. \tag{2}$$

Since each atom interacts with the cavity only once, the state of the cavity after n such interactions is given by $\rho(n) = \mathcal{T}_*^n(\rho)$ which can be interpreted as a discrete time quantum Markov dynamics.

Let Λ_n denote the random variable equal to the number of atoms detected in the ground state after n interactions. The *moment generating function* (MGF) of Λ_n is

$$\mathbb{E}\left(e^{s\Lambda_n}\right) = \sum_{k\geqslant 0} \mathbb{P}\left(\Lambda_n = k\right) e^{sk} \tag{3}$$

where the expectation is taken with respect to the probability distribution of the counting process. Using (1) it is easy to see that the MGF is equal to

$$\mathbb{E}\left(e^{s\Lambda_n}\right) = \mathrm{Tr}\left(\mathcal{T}_{*s}^n(\rho)\right) = \mathrm{Tr}\left(\rho\mathcal{T}_s^n(\mathbf{1})\right)$$

where ρ is the inital state of the cavity and $\mathcal{T}_{*s}$ (and its dual $\mathcal{T}_s$) is obtained by biasing the second Kraus operator in Eq. (1) by a factor e^s:

$$\mathcal{T}_{*s}(\rho) = L_1\rho L_1^* + e^s L_2\rho L_2^*.$$

As we will see, this allows us to study the probabilistic properties of the counting process $\Lambda(t)$ in terms of spectral properties of the operator T_s. Before that, we will make the model more realistic by passing to a continuous time description in which the atoms are Poisson distributed in time with intensity N_{ex}, and the cavity is in contact with a thermal bath. If one ignores the details of short term cavity evolution, the discrete time dynamics can be replaced by coarse grained continuous time *Lindblad (master) equation*

$$\begin{aligned} &\frac{d}{dt}\rho(t) = \mathcal{L}_*(\rho(t)),\\ &\mathcal{L}_*(\rho) = \sum_{i=1}^{4}\left(L_i\rho L_i^* - \frac{1}{2}\{L_i^*L_i, \rho\}\right). \end{aligned} \tag{4}$$

with jump operators L_i defined by

$$L_1 = \sqrt{N_{\mathrm{ex}}}\cos(\phi\sqrt{aa^*}), \tag{5}$$

$$L_2 = \sqrt{N_{\mathrm{ex}}}a^*\frac{\sin(\phi\sqrt{aa^*})}{\sqrt{aa^*}}, \tag{6}$$

$$L_3 = \sqrt{\nu+1}a, \tag{7}$$

$$L_4 = \sqrt{\nu}a^*. \tag{8}$$

As before, the operators L_1 and L_2 are associated to the detection of an atom in the excited, and respectively ground state. The emission and absorption of photons due to the contact with the bath is represented by the operators L_3 and respectively L_4. The Heisenberg picture Lindbladian $\mathcal{L}$ is the generator of a strongly continuous semigroup (cf. [13] for an introduction to the general theory) on $\mathcal{B}(\mathfrak{h})$. This means that there exists a family $(\mathcal{T}(t))_{t\geqslant 0}$ of maps on $\mathcal{B}(\mathfrak{h})$ which satisfy the *semigroup property*

$$\begin{aligned} \mathcal{T}(t)\mathcal{T}(s) &= \mathcal{T}(t+s) \quad \text{for all } t, s \geqslant 0, \\ \mathcal{T}(0) &= I, \end{aligned}$$

such that $t \mapsto \mathcal{T}(t)(X)$ is norm continuous for all $X \in \mathcal{B}(\mathfrak{h})$. The generator $\mathcal{L}$ can be recovered by

$$\mathcal{L}(X) = \lim_{h\downarrow 0} \tfrac{1}{h} \left(\mathcal{T}(h)(X) - X\right),$$

for all X in the domain of $\mathcal{L}$. Although no simple expression exists for the operators $\mathcal{T}(t)$ in terms of the generator $\mathcal{L}$, it can be helpful to think of $T(t)$ as the exponential of the generator

$$\mathcal{T}(t)(X) = e^{t\mathcal{L}}(X), \tag{9}$$

especially from the point of view of relating spectral properties of $\mathcal{L}$ to those of $\mathcal{T}(t)$, e.g. *spectral mapping theorems*. By definition, equation (9) does hold for analytic vectors of $\mathcal{L}$ which form a core of its domain.

In the discrete-time model we were able to express the moment generating function in terms of a biased transition operator. In the continuous-time case, a similar expression can be obtained in terms of a biased generator $\mathcal{L}_s$, where the jump operator L_2 associated to the ground state counting process is multiplied by the factor e^s

$$\mathcal{L}_s(X) = e^s L_1^* X L_1 - \frac{1}{2}\{L_1^* L_1, X\} + \sum_{i=2}^{4} \left(L_i^* X L_i - \frac{1}{2}\{L_i^* L_i, x\} \right). \tag{10}$$

Note that $\mathcal{L}_s$ is still the generator of a completely positive semigroup, but it is no longer identity preserving, and therefore does not represent a physical evolution except for $s = 0$. Its importance lies in the fact that the moment generating function associated to Λ_t can be expressed in analogy to equation (3) in the discrete time setting, as

$$\mathbb{E}\left(e^{s\Lambda_t}\right) = \mathrm{Tr}\left(\rho e^{t\mathcal{L}_s}(\mathbf{1})\right). \tag{11}$$

This equation plays a central role in this paper; we will use it to formulate a large deviations principle for the counting process Λ_t which analyse the asymptotic (long-time) behaviour of the process. In particular, we will use Eq. (11) to relate the moment generating function of Λ_t to the spectral properties of $\mathcal{L}_s$.

In asymptotics it is important to verify whether the Markov semigroup is ergodic, i.e. there exists a unique stationary state ρ_{ss} (i.e. $\mathcal{L}_*(\rho_{ss}) = 0$), such that

$$\lim_{t\to\infty} \mathcal{T}_*(t)(\rho) = \rho_{ss},$$

for all initial states ρ. In the case of the atom maser, such a stationary state exists and is unique [14]. Since the semigroup generated by $\mathcal{L}$ leaves invariant the commutative subalgebra generated by the number operator N, the stationary state must be diagonal in the number basis. The restriction of $\mathcal{T}(t)$ diagonal algebra we obtain a classical process known as a *birth-death process*, for which the (unique) stationary state can be computed explicitly as

$$\rho_{ss}(n) :== \rho_{ss}(0) \prod_{k=1}^{n} \left(\frac{\nu}{\nu+1} + \frac{N_{ex}}{\nu+1} \frac{\sin^2(\phi\sqrt{k})}{k} \right). \tag{12}$$

where $\rho_{ss}(n)$ is the probability of finding n photons in the cavity and $\rho_{ss}(0)$ is taken such that $\mathrm{Tr}(\rho_{ss}) = 1$. The *mean photon number* of the stationary state is given by

$$\langle \mathsf{N} \rangle = \sum_{n \geqslant 0} n \rho_{ss}(n). \tag{13}$$

The dependence of the mean photon number and photon number distribution on the parameter ϕ is shown in Fig. 1, with the values $\nu = 0.15$ and $N_{ex} = 100$ for the other parameters. We note two interesting features in this figure: first, there is a sharp change in the mean photon number at $\alpha = 1$ followed by less pronounced jumps near $\alpha = 2\pi$ and $\alpha = 4\pi$.

The other, related, feature to note concerns the photon number distribution, which only has a single peak for most values of α. In certain critical regions for α, such as $4.6 < \alpha < 7.8$, this distribution becomes bimodal, indicating that the stationary state is *bistable*. Such *metastable* behaviour of the atom maser has been investigated, using different techniques, in Ref. [15]. We study this metastable behaviour using the large deviations formalism. The large deviations rate function, introduced below, can be used as an indicator of the sharp jumps in the mean photon number.

Large deviations

The main result of this paper is the existence of what is called a *large deviations principle* for the counting process Λ_t introduced above. In [1,2] the stationary dynamics of the atom maser is analysed using the theories of large deviations and dynamical phase transitions. The large deviations approach raises important questions related to the existence of dynamical phase transitions which can be formulated in terms of the spectral properties of the perturbed generator $\mathcal{L}_s$ defined above. We note that a large deviations principle has been shown to hold in [7] in the context of quantum spin chains; the main difference is that we are looking at a continuous time Markov process on an infinite-dimensional space.

We follow [1, 2] in using large deviations to analyse quantum dynamical phase transitions, where a biased generator is interpreted as creating trajectories with a greater or lower number of ground state atoms emerging from the atom maser. This biased generator appears in Eq. 10 where we use it to define the moment generating function associated to the counting process Λ_t.



Figure 1: Mean photon number (white line) and photon number distribution (background) in the stationary state ρ_{ss} as function of $\alpha = \sqrt{N_{ex}}\phi$

This biased generator $\mathcal{L}_s$, although it is not physical due to not being trace preserving, changes the expected number of atom jumps from excited to ground state, depending on the sign of s.

We will here briefly explain what we mean by a large deviations principle and introduce some of the underlying theory. We refer the reader to [16] for a complete reference; see [17] for a comprehensive overview and [18] for an introduction to large deviations in the context of statistical mechanics. Large deviations is a framework for studying certain types of behaviour of random processes (that is, sequences of random variables). In particular, it serves to quantify what very rare events are, and to analyse the probabilities of such events. Asymptotically (in the sense of the sequence of random variables), such rare events will have a probability which decays exponentially, with the rate of exponential decay given by a *rate function*. One of the main results, which we will use, is the *Gärtner-Ellis theorem*, which relates this rate function to the limiting behaviour of the moment generating functions associated to the random variables.

Since a sequence of random variables is entirely characterised by their respective probability distributions (or laws), the following results are all formulated in terms of a sequence of probability measures. In particular, the *large deviation principle* (LDP) will characterise the limiting behaviour, as $n \to \infty$, of a family of probability measures $(\mu_n)_{n\in\mathbb{N}}$ on some topological space $(\mathcal{X}, \mathcal{B})$ in terms of a rate function. A function $I : \mathcal{X} \to [0, \infty]$ is called a *rate function* if it is lower semicontinuous (that is, its level sets $\{x \in \mathcal{X} : I(x) \leqslant \alpha\}$ are closed); if in addition its level sets are compact, we call I a *good rate function*. The domain of I is the set of points in $\mathcal{X}$ for which I is finite.



Figure 2: $\lambda'(s)$ and the spectral gap of L_s as functions of s and $\alpha = \phi/\sqrt{N_{ex}}$

The informal idea behind the large deviations principle is the exponential decay of the probabilities assigned by the family $(\mu_n)_{n\in\mathbb{N}}$ to events $x \in \mathcal{X}$, which we may illustrate by writing

$$\mu_n(dx) \approx e^{-nI(x)}dx$$

where I is the rate function associated to the family of probability measures. We will now state the large deviations principle rigorously. The limiting behaviour of the probability measures $\{\mu_n\}$ is characterised in terms of asymptotic upper and lower bounds on the values that μ_n assigns to measurable subsets $\Gamma \in \mathcal{B}$. The sequence of probability measures $\{\mu_n\}$ satisfies a large deviation principle with a rate function I if, for all $\Gamma \in \mathcal{B}$,

$$-\inf_{x\in\Gamma^0} I(x) \leqslant \liminf_{n\to\infty} \tfrac{1}{n} \log \mu_n(\Gamma) \leqslant \limsup_{n\to\infty} \tfrac{1}{n} \log \mu_n(\Gamma) \leqslant -\inf_{x\in\bar{\Gamma}} I(x).$$

The sentence "μ_n satisfies the LDP" is used as shorthand for "$\{\mu_n\}$ satisfies the large deviation principle with rate function I."

We want to prove the LDP for the counting process Λ_t of the atom maser; we will do this not by showing that Λ_t satisfies the above definition directly, but by applying the Gärtner-Ellis theorem to Λ_t. This theorem gives sufficient conditions on the sequence of probability measures in order to satisfy an LDP; its utility lies in its generality, as it does not require the probability measures to be independent.

Theorem 1 (Gärtner-Ellis theorem [16], **pp**. 44-55)**.** Let (Z_n) be a sequence of random variables in $\mathbb{R}^d$ with laws μ_n. Suppose that the logarithmic moment generating function

$$\Lambda(\lambda) = \lim_{n\to\infty} \tfrac{1}{n} \log \mathbb{E}\left[e^{\langle n\lambda, Z_n\rangle}\right], \quad \lambda \in \mathbb{R}^d$$

exists as an extended real number and is finite in a neighbourhood of the origin, and let Λ^* denote the Fenchel-Legendre transform of Λ, given by

$$\Lambda^*(x) = \sup_{\lambda \in \mathbb{R}^d} \{\langle \lambda, x \rangle - \Lambda(\lambda)\}.$$

If Λ is an essentially smooth, lower semicontinuous function (e.g. Λ is differentiable on $\mathbb{R}^d$) then (Z_n) satisfies a LDP with good rate function Λ^*.

The large deviation approach involves the use of the perturbed generator introduced in Eq. (10), and its connection to the moment generating function of the process Λ_t given by Eq. (11). In this setting, the Gärtner-Ellis theorem tells us that the sequence of atom counts Λ_t satisfies a large deviation principle, meaning that

$$\lim_{t\to\infty} \tfrac{1}{t} \log \mathbb{P}\left(\Lambda_t \leqslant a\right) = I(a),$$

where $I(a)$ is the Legendre transform of the function $\lambda(s)$ defined by

$$\lambda(s) = \lim_{t\to\infty} \frac{1}{t} \log \mathbb{E}\left(e^{s\Lambda_t}\right), \tag{14}$$

whenever this function exists and is differentiable. Therefore, to prove the desired large deviation result we show that this function $\omega_0(s)$ indeed satisfies these conditions. We will do this by showing that $\omega_0(s)$ is exactly the *spectral bound* of the generator $\mathcal{L}_s$, that is, the largest real part of the eigenvalues of $\mathcal{L}_s$. An intermediate step is relating $\omega_0(s)$ to the *spectral radius* of the operators $\exp(t\mathcal{L}_s)$ in the semigroup generated by $\mathcal{L}_s$, which involves using a Krein-Rutman theorem to show that this spectral radius is an eigenvalue of $\exp(t\mathcal{L}_s)$. We will also show that $\omega_0(s)$ is not only differentiable as a function of s, but even analytic. This serves to prove that there always exists a non-zero *spectral gap* between ω_s and the rest of the spectrum of $\mathcal{L}_s$; this spectral analysis is illustrated in Fig.2. Before we go on to state and prove our main result, we need to introduce a few more concepts from operator algebra, including the aforementioned Krein-Rutman theorem.

Spectral properties of operators

One of the main results we rely on in proving this large deviations result is the Krein-Rutman theorem. In the context of operators on a Banach space, this says that the spectral radius of an operator is an eigenvalue with a positive eigenvector, so long as the operator satisfies certain properties. Before we state the Krein-Rutman theorem, we introduce some concepts from operator algebra; for details we refer the reader to [19].

We are considering a semigroup of maps on the (infinite-dimensional) Banach space $\mathcal{B}(\mathfrak{h})$ generated by $\mathcal{L}_s$ defined in Eq. (10). In our proof we rely on spectral properties of such maps. Recall that the *spectral radius* $r(T)$ of an operator T on a Banach space X is given by

$$r(T) = \sup\{|\lambda| : \lambda \in \sigma(T)\}$$

where $\sigma(T)$ denotes the spectrum of T. The operator T is called *compact* if the image of any bounded subset of X is a relatively compact subset; however, we will not use this definition directly, and will rely on sufficient conditions to obtain compactness of the semigroup. The spectrum of a compact operator is always a countable set containing 0, that is, if T is compact then

$$\sigma(T) = \{0\} \cup \{\lambda_n : n \in \mathbb{N}\}$$

where $\lim_{n\to\infty} = 0$. The *resolvent operator* $R(z, T)$ is an important tool used to relate the spectral properties of the generator L_s to that of an element in the semigroup; it is defined by

$$R(z, T) = (T - \rho)^{-1} \quad \text{for all } z \in \mathbb{C} \setminus \sigma(T).$$

In particular, it is used to establish a *spectral mapping theorem* (see [20]) which gives the spectrum of $\exp(t\mathcal{L}_s)$ in terms of the spectrum of $\mathcal{L}_s$,

$$\sigma(e^{tL_s}) = \left\{ e^{t\lambda} : \lambda \in \sigma(L_s) \right\}.$$

For this we rely on compactness of the resolvent $R(z, \mathcal{L}_s)$, which is a property we will prove below. The Krein-Rutman theorem gives more precise information about the spectral radius of a compact operator.

We want to apply a relevant version of the Krein-Rutman theorem to the semigroup generated by $\mathcal{L}_s$. Generally such statements are concerned with operators which preserve the positive cone in a Banach algebra; in particular, to obtain uniqueness of the spectral radius as being the only eigenvector of $\mathcal{T}_s(t)$ we need the semigroup to be strongly positive (or positivity improving). This means that, given any positive element $x \in L^2(\mathfrak{h})$, we have that $\mathcal{T}_s(t)(x)$ is strictly positive (that is, lies within the interior of the cone of positive elements).

Theorem 2 (Krein-Rutman [21], Ch. 1)**.** Let X be a Banach space, $K \subset X$ a soid cone and $T : X \to X$ a compact linear operator which is strongly positive, that is, $Tx \in \text{int}(K)$ if $x \in K \setminus \{0\}$. Then

1. The spectral radius $r(T)$ is positive and $r(T)$ is a simple eigenvalue with an eigenvector $x \in \text{int}(K)$; there is no other eigenvalue with a positive eigenvector;
2. For any other eigenvalue $\lambda \neq r(T)$ we have $|\lambda| < r(T)$.

We want to use this theorem to establish that for each $t > 0$, the spectral radius of $\mathcal{T}_s(t)$ is a simple eigenvalue and $\mathcal{T}_s(t)$ has no eigenvalues with the same modulus.

3 Results and Methodology

Our main result can be summarised as follows.

Theorem 3. The counting process Λ_t satisfies a large deviations principle.

This means that, although the behaviour of the spectral bound $\lambda(s)$ is certainly interesting (as we will discuss below), there are no non-analyticities; therefore, there are no quantum dynamical phase transitions in this sense.

The main result is based on showing a Gärtner-Ellis theorem for the counting process Λ_t. This consists of proving a Perron-Frobenius type result for the transition operators associated to the modified generator $\mathcal{L}_s$, that is, that the spectral radius $r_s(t)$ is an eigenvalue of the operator $e^{t\mathcal{L}_s}$, and no other eigenvalue has the same modulus. The remainder consists of showing that the rate function $r_s(t)$ is an analytic function of s. The analysis of the spectral properties of the operator $\mathcal{L}_s$ involves choosing a suitable representation for the action of the generator, which allows us to show that the semigroup it generates is immediately compact.

The steps in the proof are outlined as follows:

1. Define the L^2-embedding, following [10]; the semigroup $(\mathcal{T}_s(t))_{t\geqslant 0}$ gives rise to a semigroup $(T_s(t))_{t\geqslant 0}$ on the space of Hilbert-Schmidt operators. This new semigroup is used to compute the desired expectation values, and we will focus on its spectral properties.

2. Restrict the associated semigroup $(T_s(t))_{t\geqslant 0}$ to the commutative subalgebra (of multiplication operators) generated by the number operator N; this defines the generator L_s acting on $l^\infty(\mathbb{N})$.

3. Show that, in this representation L is self-adjoint and the semigroup generated by L is immediately compact.

4. Use bounded perturbation theory to show that the semigroup generated by L_s is compact.

5. Show that the semigroup generated by L_s is strictly positive.

6. Krein-Rutman theorem: the spectral radius $r_s(t)$ of $T_s(t)$ is an eigenvalue with strictly positive eigenvector.

7. Analyticity of perturbation implies $\lambda(s)$ is analytic.

Since the transition operator $T_s(t)$ is compact for all $t > 0$, it has discrete spectrum; in particular, there is a spectral gap between the top eigenvalue $r_s(t)$ and the next eigenvalue. Along with the strict positivity of the eigenvector associated to $r_s(t)$, we can prove the desired convergence of the moment generating functions; analyticity (in particular differentiability) of the rate function completes the large deviations principle.

The first step in the proof is to define a semigroup $(T_s(t))_{t\geqslant 0}$ associated to $(\mathcal{T}_s(t))_{t\geqslant 0}$; this follows [10] where this is used to find a representation of quantum dynamical semigroups. We emphasise that we do not strictly consider this new semigroup as a representation of $(\mathcal{T}_s(t))_{t\geqslant 0}$; rather, it is a separate entity $(T_s(t))_{t\geqslant 0}$ whose expectation values and asymptotic properties coincide. We define the new semigroup on the space $L^2(\mathfrak{h})$ of Hilbert-Schmidt operators on the Hilbert space $\mathfrak{h}$, which is particularly useful since on this space the generator $\mathcal{L}_0$ is self-adjoint. This means that we can consider L_s as a perturbation of a self-adjoint operator, which makes the subsequent analysis significantly easier. It is in this setting that we will work for the remainder; we simply refer to L_s for the associated generator.

This semigroup $(T_s(t))_{t \geqslant 0}$ leaves the commutative subalgebra generated by the number operator N invariant. In the remainder of this article we will consider the restriction of the semigroup to this *diagonal algebra*; by $T_s(t)$ and L_s we will implicitly mean the restrictions to this subalgebra.

As our aim is to apply the Krein-Rutman theorem to the operators $T_s(t)$, we show that these operators are compact and strictly positive. We show that the semigroup generated by L_s is an immediately compact semigroup. In order to show this, we first establish that the generator L_0 has compact resolvent and generates a semigroup of compact operators. Then, upon showing that the perturbation term in $L_s = L + \Delta_s$ is bounded, we apply bounded perturbation theory to obtain the desired result.

Because $T_s(t)$ is compact, we can apply the Krein-Rutman theorem for compact operators to claim that the spectral radius $r_s(t) = \exp(t\lambda(s))$ is an eigenvalue of $T_s(t)$, which is associated to a positive eigenvector $|R_s\rangle$. Furthermore, because $T_s(t)$ is positivity improving, the eigenvector $|R_s\rangle$ is strictly positive. This then allows us to conclude that the convergence required for the Gärtner-Ellis theorem holds, that is,

$$\lim_{t \to \infty} \frac{1}{t} \log \mathbb{E}\left(e^{s\Lambda_t}\right) = \lambda(s)$$

which establishes the large deviation principle. Finally, because L_s is an analytic perturbation of L, the function $\lambda(s)$ is itself analytic.

4 Details of proof

The proof of our main result starts with a study of the spectral properties of the perturbed generator $\mathcal{L}_s$. Before we begin, we define an associated semigroup of the dynamics by embedding the algebra $\mathcal{B}(\mathfrak{h})$ in the Hilbert space $L^2(\mathfrak{h})$ of *Hilbert-Schmidt operators* (we adapt this technique from [10]). We will refer to this associated semigroup as an *embedding* of the original semigroup.

Recall that $X \in \mathcal{B}(\mathfrak{h})$ is called a Hilbert-Schmidt operator if it has finite Hilbert-Schmidt norm, that is,

$$\|X\|^2_{\text{H-S}} = \text{Tr}|X|^2 < \infty.$$

The space of all Hilbert-Schmidt operators, denoted $L^2(\mathfrak{h})$, is a Hilbert space with inner product

$$\langle X, Y \rangle_{\text{H-S}} = \text{Tr}\,(X^*Y) \quad \text{for all } X, Y \in L^2(\mathfrak{h}). \tag{15}$$

This map which embeds $\mathcal{B}(\mathfrak{h})$ into $L^2(\mathfrak{h})$ is defined in terms of the stationary state ρ_{ss} (see Eq. (12)) of the generator $\mathcal{L}$.

Following [10], we define the embedding $\mathfrak{i}$ as

$$\begin{aligned} \mathfrak{i} &: \mathcal{B}(\mathfrak{h}) \to L^2(\mathfrak{h}) \\ &: x \mapsto \rho_{ss}^{1/4} x \rho_{ss}^{1/4}. \end{aligned}$$

Using this map $\mathfrak{i}$, we can define the embedding of operators on $\mathcal{B}(\mathfrak{h})$; with an operator $\mathcal{T} : \mathcal{D}(\mathcal{T}) \to \mathcal{B}(\mathfrak{h})$ we associate an operator T on $L^2(\mathfrak{h})$ by

$$T(\mathfrak{i}(X)) = \mathfrak{i}(\mathcal{T}(X)) \quad \text{for all } X \in \mathcal{D}(\mathcal{T}) \subset \mathcal{B}(\mathfrak{h}).$$

This embedding is used to define the associated semigroup; we formally define the generator L_s on $L^2(\mathfrak{h})$ by

$$L_s(X) = \mathfrak{i}(\mathcal{L}_s(\mathfrak{i}^{-1}(X))). \tag{16}$$

It will be helpful to write L_s in the perturbative form

$$L_s(X) = L(X) + \Delta_s(X) \tag{17}$$

where, recalling Eq. 10, L is the generator obtained by applying the embedding to the original generator $\mathcal{L}$ and Δ_s is the perturbation arising from biasing the generator. By restricting to finite-rank elements of $L^2(\mathfrak{h})$ it is impossible to find an explicit expression for the action of L_s.

We will be using this associated semigroup $(T_s(t))_{t \geqslant 0}$ in the remainder of this paper. To prove the large deviations principle for the counting process Λ_t, we express the expectation value in Eq. (11) in terms of an expectation value on $L^2(\mathfrak{h})$. Using Eq. (16) we are able to show that

$$\mathrm{Tr}\left(\rho_{in} \mathcal{T}_s(t)(\mathbf{1})\right) = \mathrm{Tr}\left(\hat{\rho}_{in} T_s(t)(\rho_{ss}^{1/2})\right)$$

where we have defined $\hat{\rho}_{in} = \rho_{ss}^{-1/4} \rho_{in} \rho_{ss}^{-1/4}$; in terms of the inner product on $L^2(\mathfrak{h})$ defined in Eq. (15), this expectation value reads

$$\mathrm{Tr}\left(\rho_{in} \mathcal{T}_s(t)(\mathbf{1})\right) = \left\langle \hat{\rho}_{in}, T_s(t)(\rho_{ss}^{1/2}) \right\rangle_{\text{H-S}}.$$

Here the action of the semigroup generated by L_s is still taken in the Heisenberg picture.

The semigroup $(T_s(t))_{t \geqslant 0}$ leaves the commutative subalgebra of multiplication operators invariant; in particular,f the stationary state ρ_{ss} is a multiplication operator. In the remainder of this paper we will only be considering the action of L_s on the commutative subalgebra $\mathcal{N}$ of $L^2(\mathfrak{h})$ consisting of multiplication operators.

In particular, for any $f \in \mathcal{D}(L_s)$ we have

$$L_s(M_f) = M_g$$

where $g \in l^2(\mathbb{N})$ is given by

$$g(n) = \lambda_{n-1}\mu_n f(n-1) - (\lambda_n^2 + \mu_n^2) f(n) + \lambda_n \mu_{n+1} f(n+1) \tag{18}$$
$$+ (e^{-1} - 1)\frac{\mu_{n+1}}{\lambda_n} N_{\text{ex}} \sin^2(\phi\sqrt{n+1}) f(n+1). \tag{19}$$

Here the *rates* λ_k, μ_k are given by

$$\lambda_k = \sqrt{\nu(k+1) + N_{\text{ex}} \sin^2(\phi\sqrt{k+1})},$$
$$\mu_k = \sqrt{(\nu+1)k}.$$

In the unperturbed case $s = 0$, Eq. (18) shows that the restriction of L_s to the commutative subalgebra of $L^2(\mathfrak{h})$ generated by $l^2(\mathbb{N})$ corresponds to a *birth-death process* from classical probability theory with rates λ_k, μ_k. Therefore L coincides with the generator of this classical birth-death process, where for any $f \in \mathcal{D}(L)$

$$L(f)(n) = \lambda_{n-1}\mu_n f(n-1) - (\lambda_n^2 + \mu_n^2) f(n) + \lambda_n \mu_{n+1} f(n+1). \tag{20}$$

The perturbation in Eq. (17) takes the following explicit form: for any $f \in \mathcal{D}(L)$

$$\Delta_s(f)(n) = (e^{-1} - 1)\frac{\mu_{n+1}}{\lambda_n} N_{\text{ex}} \sin^2(\phi\sqrt{n+1}) f(n+1). \tag{21}$$

This defines the representation L_s on the Hilbert space $l^2(\mathbb{N})$ of the perturbed associated generator; abusing notation, we clearly have

$$L_s(M_f) = M_{L_s(f)} \quad \text{for all } f \in \mathcal{D}(L).$$

The remainder of the proof will focus on the spectral properties of the generator L_s.

Lemma 1. The semigroup generated by L is immediately compact.

Proof. Since the L^2-embedding defines L as a self-adjoint operator on the Hilbert space $l^2(\mathbb{N})$, the spectral theorem applies. Let the spectral decomposition of L be of the form

$$L = \sum_{n \geqslant 0} \lambda_n P_n,$$

where $\sigma(L) = (\lambda_n)_{n\geqslant 0}$ is the spectrum of L and P_n is the eigenprojection associated to λ_n. Then the spectral theorem for self-adjoint operators states that

$$T_t = \sum_{n \geqslant 0} e^{t\lambda_n} P_n \quad \text{for all } t \geqslant 0. \tag{22}$$

This ansatz will be used to show that T_t is compact for every $t > 0$; in particular, we will show that T_t may be approximated in norm by the partial sums in Eq. (22). In order for this to be valid, we show that L has discrete spectrum (in the sense of having a countable number of eigenvalues and finite-dimensional eigenspaces), and the eigenvalues of L diverge to $-\infty$.

In order to establish that L has discrete spectrum, we follow the argument used in [22]. Note that the action (see Eq. (20)) of L on $l^2(\mathbb{N})$ may be represented by the infinite-dimensional tri-diagonal matrix

$$L = \begin{pmatrix} -(\lambda_0^2+\mu_0^2) & \lambda_0\mu_1 & 0 & \\ \lambda_0\mu_1 & -(\lambda_1^2+\mu_1^2) & \lambda_1\mu_2 & \\ 0 & \lambda_1\mu_2 & -(\lambda_2^2+\mu_2^2) & \\ & & & \ddots \end{pmatrix} \tag{23}$$

acting on infinite column vectors representing $f \in l^2(\mathbb{N})$. We split L up into a sum

$$L = A + B$$

of two infinite-dimensional matrices; A containing the diagonal elements in Eq. (23), and B containing the off-diagonal elements. Then L can be regarded as a perturbation of A; in particular, the perturbation is relatively bounded in the sense of Kato [23], that is,

$$\|Bx\| \leqslant \|Ax\| \quad \text{for all } x \in \mathcal{D}(A).$$

The resolvent of A is compact, and using Kato's result on relatively bounded perturbations (p. 214) we find that the resolvent of $L = A + B$ is also compact. Since the self-adjoint operator L is closed (see [24], p. 256), it follows (Kato, p. 187) that the spectrum of L is discrete.

Consider the the partial sums of the exponential of the generator L defined by

$$T^{(N)}(t) = \sum_{n=0}^{N} e^{t\lambda_n} P_n,$$

where $\{\lambda_1, \lambda_2, \ldots\}$ is the spectrum of L and P_n is the projection onto the eigenspace corresponding to λ_n. The real eigenvalues $\{\lambda_1, \lambda_2, \ldots\}$ are bounded from above and by using the spectral mapping theorem on the spectrum of the compact resolvent of L we conclude that $\lambda_n \to -\infty$. Therefore e^{tL} may be approximated by the finite-rank partial sums and e^{tL} is compact for all $t > 0$. □

Having established that the semigroup generated by L is compact, we use a bounded perturbation argument to extend this property to the semigroup generated by L_s.

Corollary 1. The semigroup generated by L_s is immediately compact.

Proof. Recall that the perturbation in Eq. (17) may be represented on the commutative subalgebra $l^2(\mathbb{N})$ as

$$L_s = L + \Delta_s$$

where Δ_s is defined in Eq. (21). Since Δ_s is a bounded operator, the semigroup generated by L_s is also immediately compact (see [13], p. 167). □

The Krein-Rutman theorem also requires that the operators $T_s(t)$ are all strictly positive, that is, $\langle e_n | T_s(t)(\rho) | e_n \rangle > 0$ for all $n \geqslant 0$. This follows from a Dyson expansion (see [25]) applied to the generator $\mathcal{L}_s = \mathcal{L}^0 + \mathcal{L}^s$, where $\mathcal{L}^0$ is the generator obtained from $\mathcal{L}_s$ by taking

$$\mathcal{L}_s(\rho) = \mathcal{L}^0(\rho) + e^{-s} L_1 \rho L_1^* + L_2 \rho L_2^*.$$

The resulting generator $\mathcal{L}^0$ is associated to a classical birth-death process and so the commutative restriction is irreducible, and so strictly positive. As the leading term in the Dyson expansion involves only $\mathcal{L}^0$, and all other terms (involving the jump operators L_1, L_2) are positive, this suffices to conclude that the operators $T_s(t)$ are strictly positive.

Our goal is to show that the spectral bound $\lambda(s)$, defined as the largest real part of the spectrum

$$\lambda(s) = \{\mathrm{Re}(\lambda) : \lambda \in \sigma(L_s)\}$$

plays the role of the large deviations rate function as introduced in Eqs. (11),(14). The first step in establishing this follows from compactness of the operators $T_s(t)$ and the Krein-Rutman theorem.

Corollary 2. The spectral radius $r_s(t)$ of $T_s(t)$ is an eigenvalue of $T_s(t)$ and has associated to it a strictly positive eigenvector.

Proof. Compactness of the semigroup $(T_s(t))_{t \geqslant 0}$ leads ([13], p. 281) to a spectral mapping theorem of the form

$$e^{t\sigma(L_s)} = \sigma(T_s(t)) \setminus \{0\} \quad \text{for all } t \geqslant 0.$$

Since each operator $T_s(t)$ is compact and strictly positive, we can apply the Krein-Rutman theorem for compact operators (see Thm. 2 above). □

Since $r_s(t) \in \sigma(T_s(t))$, using the spectral mapping theorem this means that

$$r_s(t) = e^{t\lambda(s)}$$

where $\lambda(s)$ is the spectral bound of L_s; any other eigenvalue $\lambda \in \sigma(T_s(t))$ satisfies $|\lambda| < r_s(t)$.

To complete the proof of the large deviations principle, we need to show that the spectral radius $r_s(t)$ of $T_s(t)$ is a differentiable function of s. Using analytic perturbation theory for the generator L_s, we can in fact prove that the spectral bound $s(L_s)$ is an analytic function of s.

The results used on analytic perturbation theory are found in [26]. Given a closed operator T' we define the family of perturbations $T(s)$ by

$$T(s) = T' - sH,$$

where H is a closed operator with the same domain as T'. The main result (Prop. 3.25, p. 141) on analycity is that any isolated eigenvalue (of finite multiplicity) and its associated eigenprojection are analytic functions of s in some disc around $s = 0$. We apply this result to the family of perturbations

$$L_s = L + \Delta_s,$$



Figure 3: Comparison of classical birth (solid) and death (dashed) rates (see text); stationary distribution

for which it is clear that the spectral bound of $\mathcal{L}_s$ is an analytic function of s. Since, by the Krein-Rutman theorem, the spectral bound of L_s and the spectral radius $r_s(t)$ are related by

$$r_s(t) = e^{s(L_s)},$$

we immediately see that $r_s(t)$ is also an analytic function of s. As a consequence of this, the spectral radius remains isolated when viewed as a function of s (which should not be confused with its isolation for any fixed s since that trivially follows from compactness of the semigroup).

5 Numerics and discussion

We have shown above that the counting process Λ_t satisfies a large deviations principle; in particular, the spectral radius $r_s(t)$ is an analytic function of the large deviations parameter s. We will now briefly consider the interpretation of the large deviations rate function in the context of quantum dynamical phase transitions.

In [1] the interpretation of the large deviations rate function as an indicator of quantum dynamical phase transitions. Non-analyticities in the rate function, such as discontinuities in the first derivative correspond to the spectral gap becoming very small (see Fig. 2). At these values for α we also see that the stationary state becomes bistable; that is, instead of a single peak in the mean photon number distribution, we see two separated peaks (see Fig. 1). This means that the state of the cavity can be in either of two bistable phases, one more active than the other; these two phases become apparent in simulations. The number of excitations in the cavity spends some time in a more active region before making a transition to a less active region. The occurrence of this behaviour for some values of α signifies a different dynamical region from other values of α, and this is a quantum dynamical phase transition.

Now consider the jumps in the mean photon number (solid line in Fig. 3). The values for α associated to the midpoint of the jump (e.g. $\alpha_c \approx 6.7$) are associated to points where the derivative $\lambda'(s)$ of the spectral bound λ crosses the s-axis in the following sense. The derivative $\lambda'(s)$ shows clearly different



(a) $\lambda'(s)$ (as in Fig. 2)

(b) Crossing at α_c for $N_{ex} = 20, 30, 90, 100$ (varying s axes)

(c) Mean photon number (see text)

Figure 4: Analysis of crossing near α_c

values when comparing either sides of the 'phase lines' visible in Fig. 4 (a). Although this figure suggests that this phase line approaches the $s = 0$ axis strictly from $s < 0$, Fig. 4 (b) shows that this line crosses over to the $s > 0$ region. The angle at which the crossing occurs increases with N_{ex} (note the changing s ranges: the top two figures are with $|s| \leqslant 0.01$ while the bottom two figures are with $|s| \leqslant 10^{-8}$).

We are interested in the $s = 0$ behaviour since, if $\lambda'(0)$ is discontinuous, this means there is a dynamical phase transition: the mean $\langle \Lambda_t \rangle$ (first moments) depends on whether the derivative is taken from the left or right. Since the phase line does cross the $s = 0$ axis, this does not necessarily preclude such a phase transition. However, we know mathematically that the spectral bound λ is an analytic function of s, so a phase transition cannot occur.

Accuracy of the numerical derivative of the rate function is illustrated in Fig. 4 (c). If the derivative with respect to s is taken with the differential Δs small enough (10^{-8}), the derivative follows the mean photon number exactly. For larger Δs, the derivative follows the 'wrong' phase and thus results in the wrong mean being computed, which is clearly visible in the top figure.

Another way of looking at these metastable phases is by considering the classical birth-death process associated to the quantum dynamical system. Following the discussion in [27], we consider the values of α for which the 'birth' rates λ_k exceed the 'death' rates μ_k along with the dynamical nature of these crossover points. In Fig. 3 we see these crossover points; for example, at $\alpha = 1$ the rates cross at a maximum for the death rate, which in the stationary state corresponds to the sharpest increase in the mean photon number. At $\alpha = 4.6$ the rates intersect in more than one location for the first time, which is reflected in the photon number distribution as the start of the bistable region. (Note that this figure is independent of the variable N_{ex}, as x here is an appropriate rescaling of n in the limit $N_{ex} \to \infty$.)

6 Concluding remarks

We have studied the counting process associated to photon number trajectories of the cavity state in the atom maser, and shown that this process satisfies a large deviations principle. Our approach could be extended to measurements other than the counting of ground-state atoms used here, for example, homodyne detection. This means taking a different perturbation of the generator to analyse the large deviations behaviour. However, the resulting dynamics will no longer have the nice property of leaving the diagonal subalgebra invariant, so the algebra and the numerical analysis will be more involved, requiring non-commutative analysis. The method could also be extended to any immediately compact, strictly positive semigroup; that is, the exact form of the generator does not directly affect the principles used here. Related results on a generalisation large deviations behaviour of the counting statistics in quantum spin chains (as addressed in [7]) will be discussed in a forthcoming publication.

References

[1] J. P. Garrahan, A. Armour, and I. Lesanovsky, "Quantum trajectory phase transitions in the micromaser,"*Physical Review E* **84** (Aug., 2011) , `arXiv:1103.0919`. `http://arxiv.org/abs/1103.0919http://link.aps.org/doi/10.1103/PhysRevE.84.021115`.

[2] J. P. Garrahan and I. Lesanovsky, "Thermodynamics of Quantum Jump Trajectories,"*Physical Review Letters* **104** (Nov., 2010) 5, `arXiv:0911.0556`. `http://arxiv.org/abs/0911.0556`.

[3] J. P. Garrahan, R. L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk, and F. van Wijland, "Dynamical First-Order Phase Transition in Kinetically Constrained Models of Glasses,"*Physical Review Letters* **98** (May, 2007) 0–3. `http://link.aps.org/doi/10.1103/PhysRevLett.98.195702`.

[4] L. O. Hedges, R. L. Jack, J. P. Garrahan, and D. Chandler, "Dynamic order-disorder in atomistic models of structural glass formers.,"*Science* **323** (Mar., 2009) 1309–13. `http://www.ncbi.nlm.nih.gov/pubmed/19197025`.

[5] J. P. Garrahan, R. L. Jack, V. Lecomte, K. van Duijvendijk, and E. Pitard, "First-order dynamical phase transition in models of glasses: an approach based on ensembles of histories," `arXiv:arXiv:0810.5298v2`. `http://arxiv.org/abs/0810.5298`.

[6] A. A. Budini, "Thermodynamics of quantum jump trajectories in systems driven by classical fluctuations," `arXiv:1012.0776`. `http://arxiv.org/abs/1012.0776`.

[7] F. Hiai, M. Mosonyi, and T. Ogawa, “Large deviations and Chernoff bound for certain correlated states on a spin chain,” *Journal of Mathematical Physics* **48** (2007) no. 12, 123301. `http://link.aip.org/link/JMAPAQ/v48/i12/p123301/s1&Agg=doi`.

[8] Y. Ogata, “Large Deviations in Quantum Spin Chains,”*Communications in Mathematical Physics* **296** (Feb., 2010) 35–68. `http://www.springerlink.com/index/10.1007/s00220-010-0986-y`.

[9] H. Comman, “Large Deviations for Quantum Markov Semigroups on the 2 ÃŮ 2-Matrix Algebra,”*Annales Henri Poincaré* **9** (Aug., 2008) 979–1003. `http://www.springerlink.com/index/10.1007/s00023-008-0379-3`.

[10] R. Carbone and F. Fagnola, “Exponential L2-convergence of quantum Markov semigroups on B(H),”*Mathematical Notes* **68** (Oct., 2000) 452–463. `http://www.springerlink.com/index/10.1007/BF02676724`.

[11] F. Fagnola, R. Rebolledo, and C. Saavedra, “Quantum flows associated to master equations in quantum optics,” *Journal of Mathematical Physics* **35** (1994) no. 1, 1. `http://link.aip.org/link/JMAPAQ/v35/i1/p1/s1&Agg=doi`.

[12] B.-G. Englert and G. Morigi, “Five Lectures on Dissipative Master Equations,” in *Coherent Evolution in Noisy Environments*, A. Buchleitner and K. Hornberger, eds., vol. 2 of *Lecture Notes in Physics*, ch. 2, pp. 55–106. Springer, Berlin Heidelberg New York, 2002. `http://www.springerlink.com/content/25m5025776145508/`.

[13] K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*. Graduate Text in Mathematics. Springer-Verlag, New York, June, 2000.

[14] F. Fagnola and R. Rebolledo, “On the existence of stationary states for quantum dynamical semigroups,” *Journal of Mathematical Physics* **42** (2001) no. 3, 1296. `http://link.aip.org/link/JMAPAQ/v42/i3/p1296/s1&Agg=doi`.

[15] L. Bruneau and C.-a. Pillet, “Thermal Relaxation of a QED Cavity,”*Journal of Statistical Physics* **134** (Dec., 2008) 1071–1095. `http://www.springerlink.com/index/10.1007/s10955-008-9656-2`.

[16] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*, vol. 38 of *Stochastic Modelling and Applied Probability*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2010. `http://www.springerlink.com/index/10.1007/978-3-642-03311-7`.

[17] R. S. Ellis, “An Overview of the Theory of Large Deviations and Applications to Statistical Mechanics,” *Scand. Actuarial. J.* (1995) no. 1, 97–142.

[18] H. Touchette, "The large deviation approach to statistical mechanics,"*Physics Reports* **478** (July, 2009) 1–69. `http://linkinghub.elsevier.com/retrieve/pii/S0370157309001410`.

[19] E. B. Davies, *Linear Operators and their Spectra*. Cambridge studies in advanced mathematics. Cambridge University Press, Cambridge, 2007.

[20] K.-J. Engel and R. Nagel, *A Short Course on Operator Semigroups*. Universitext. Springer, New York, 2006.

[21] Y. Du, *Order Structure And Topological Methods in Nonlinear Partial Differential Equations*, vol. 1 of *Series on Partial Differential Equations and Applications*. World Scientific, Singapore, 2006.

[22] J. Janas and S. Naboko, "Multithreshold Spectral Phase Transitions for a Class of Jacobi Matrices," in *Recent Advances in Operator Theory: The Israel Gohberg Anniversary Volume*, vol. 124 of *Operator Theory: Advances and Applications*, pp. 267–285. Birkhäuser Verlag, Basel Boston Berlin, 2001.

[23] T. Kato, *Perturbation Theory for Linear Operators*. Springer-Verlag, Berlin Heidelberg New York, 2nd ed., 1976.

[24] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, vol. I. Academic Press, New York, 1972.

[25] C.-A. Pillet, "Quantum Dynamical Systems," in *Open Quantum Systems I*, no. Umr 6207 in Lecture Notes in Mathematics, pp. 107–182. Berlin / Heidelberg, 2006. `http://www.springerlink.com/content/717852g1268mhqn3/`.

[26] F. Chatelin, *Spectral Approximation of Linear Operators*. Computer Science and Applied Mathematics. Academic Press, London, 1983.

[27] B.-G. Englert, "Elements of Micromaser Physics," `arXiv:0203052 [quant-ph]`. `http://arxiv.org/abs/quant-ph/0203052`.

[28] F. Cipriani, F. Fagnola, and J. M. Lindsay, "Spectral Analysis and Feller Property for Quantum Ornstein-Uhlenbeck Semigroups,"*Communications in Mathematical Physics* **210** (Mar., 2000) 85–105. `http://www.springerlink.com/openurl.asp?genre=article&id=doi:10.1007/s002200050773`.

A Equality of expectation values for $\mathcal{L}_s$ and L_s

Instead of considering the embedded generator L_s as a representation of L_s on $L^2(\mathfrak{h})$, we should interpret these as generators of separate semigroups, one on $L^2(\mathfrak{h})$ and the other on $\mathcal{B}(\mathfrak{h})$, and show that they agree in the expectation value we are interested in. That is, show that the quantities defined by the moment generating function

$$\mathbb{E}(e^{s\Lambda_t}) = \mathrm{Tr}(\rho_{\mathrm{in}}\mathcal{T}_s(t)(\mathbf{1}))$$

and the inner product

$$\mathrm{Tr}(\tilde{\rho}_{\mathrm{in}}e^{tL_s}(\rho^{1/2})) = \langle\tilde{\rho}_{\mathrm{in}}, T_s(t)(\rho^{1/2})\rangle_{\text{H-S}}$$

are finite, and equal, for all finite-rank initial states ρ_{in} and all $t > 0$. Note that the moment generating function should be finite since it is bounded by the moment generating function for a Poisson process

$$\mathbb{E}\left(e^{s\Lambda_t}\right) = \sum_{k\geqslant 0} e^{sk}\mathbb{P}_\rho\left(\Lambda_t = k\right) \tag{24}$$

$$\leqslant \sum_{k\geqslant 0} e^{sk}\mathbb{P}_{\mathrm{Poisson}}(k) \tag{25}$$

since the arrival times of the atoms are governed by a Poisson distribution.

Following a similar argument in [28], we show that there exists a time $T > 0$ such that for all $0 \leqslant t < T$, the series expansion

$$e^{t\mathcal{L}_s}(x) = \sum_{k\geqslant 0} \frac{t^k\mathcal{L}_s^k(x)}{k!}$$

defines a bounded operator for all finite rank operators $x \in \mathcal{B}(\mathfrak{h})$. In the language of operator analysis, all finite-rank operators are *analytic vectors* for $\mathcal{L}_s$. This is done by considering rank one operators of the form $|e_n\rangle\langle e_m|$ and finding a $T > 0$ such that the series

$$\sum_{k\geqslant 0} \frac{t^k\|\mathcal{L}_s(|e_n\rangle\langle e_m|)\|}{k!}$$

converges for all $t < T$. Such a T is readily found (see [28]) and depends on the coefficients μ, λ appearing in the master equation.

Having shown that all finite rank operators are analytic vectors for $\mathcal{L}_s$, we continue by noting that given any operator $x \in \mathcal{B}(\mathfrak{h})$ we can define its (finite rank) projection

$$x_n = P_n x P_n$$

where P_n projects onto the subspace of rank n operators. We are interested in the action of the semigroup on trace-class operators; the generator of this *predual semigroup* is denoted by $\mathcal{L}_{s*}$ and is induced by $\mathcal{L}_s$ by the duality relation

$$\mathrm{Tr}(e^{t\mathcal{L}_{s*}}(\rho)x) = \mathrm{Tr}(\rho e^{t\mathcal{L}_s}(x))$$

for all $\rho \in \mathcal{T}_1(\mathfrak{h})$ and all $x \in \mathcal{B}(\mathfrak{h})$.

We want to show equality in

$$\mathrm{Tr}(\rho_{\mathrm{in}}e^{t\mathcal{L}_s}(\mathbf{1})) = \mathrm{Tr}(\tilde{\rho}_{\mathrm{in}}e^{tL_s}(\rho^{1/2}))$$

for all $t > 0$, (for now) for an appropriately chosen set of initial states ρ_{in}. Partition the time t by writing $t = t_1 + \ldots + t_l$ where $t_i < T$. The main idea behind showing the above equality is by applying sequentially the analyticity of finite-rank operators and weak-$*$ continuity to 'move over' each element of the semigroup $e^{t_i \mathcal{L}}$ to the trace-class operators and then the Hilbert-Schmidt operators, and repeat this until the entire semigroup lives on the other side. Using the projections P_n introduced above, we find that by weak-$*$ continuity and the semigroup property we have

$$\operatorname{Tr}\left(\rho e^{(t_1+\ldots+t_l)\mathcal{L}_s}(\mathbf{1})\right) = \lim_{n\to\infty} \operatorname{Tr}\left(\rho e^{t_1\mathcal{L}_s}\left(P_n e^{(t_2+\ldots+t_l)\mathcal{L}_s}(\mathbf{1})P_n\right)\right).$$

Define the finite-rank operator $A_n = P_n e^{(t_2+\ldots+t_l)\mathcal{L}_s}(\mathbf{1})P_n$; then since this is an analytic vector by the above argument, we may express the RHS as a power series and thus we obtain

$$\lim_{n\to\infty} \operatorname{Tr}\left(\rho \sum_{k\geqslant 0} \frac{t_1^k \mathcal{L}_s^k}{k!}(A_n)\right) = \lim_{n\to\infty} \operatorname{Tr}\left(\rho_{ss}^{-1/4}\rho\rho_{ss}^{-1/4} \sum_{k\geqslant 0} \frac{t_1^k L_s^k}{k!}(\rho_{ss}^{1/4} A_n \rho_{ss}^{1/4})\right)$$

by definition of the embedded generator L_s. Write the embedded initial state as $\tilde{\rho} = \rho_{ss}^{-1/4}\rho\rho_{ss}^{-1/4}$ and this limit may be written as an inner product in the Hilbert-Schmidt space as

$$\begin{aligned}
\lim_{n\to\infty} \operatorname{Tr}\left(\tilde{\rho} e^{t_1 L_s}\left(\rho_{ss}^{1/4} P_n A P_n \rho_{ss}^{1/4}\right)\right) &= \lim_{n\to\infty} \langle \tilde{\rho}, e^{t_1 L_s}(A_n)\rangle && (26)\\
&= \lim_{n\to\infty} \langle e^{t_1 L_{s*}}(\tilde{\rho}), A_n\rangle. && (27)
\end{aligned}$$

It now suffices to show that $A_n \to A$ weakly; this is clear, since for any $\tau \in \mathcal{T}_1(\mathfrak{h})$ we have

$$\begin{aligned}
\langle \tau, P_n \rho_{ss}^{1/4} A \rho_{ss}^{1/4} P_n\rangle &= \operatorname{Tr}\left(\tau P_n \rho_{ss}^{1/4} A \rho_{ss}^{1/4} P_n\right) && (28)\\
&= \operatorname{Tr}\left(\rho_{ss}^{1/4}\tau\rho_{ss}^{1/4} P_n A P_n\right) && (29)\\
&\to \operatorname{Tr}\left(\rho_{ss}^{1/4}\tau\rho_{ss}^{1/4} A\right); && (30)
\end{aligned}$$

whenever $\rho_{ss}^{1/4}\tau\rho_{ss}^{1/4} \in \mathcal{T}_1(\mathfrak{h})$,

$$\operatorname{Tr}\left(\rho_{ss}^{1/4}\tau\rho_{ss}^{1/4}\right) = \operatorname{Tr}\left(\rho_{ss}^{1/2}\tau\right) = \langle \rho_{ss}^{1/2}, \tau\rangle < \infty$$

so

$$P_n \rho_{ss}^{1/4} A \rho_{ss}^{1/4} P_n \to \rho_{ss}^{1/4} A \rho_{ss}^{1/4}$$

weakly. Therefore the limit becomes

$$\begin{aligned}
&\operatorname{Tr}\left(\tilde{\rho} e^{t_1 L_s}\left(\rho_{ss}^{1/4} e^{(t_2+\ldots+t_l)\mathcal{L}_s}(\mathbf{1})\rho_{ss}^{1/4}\right)\right) && (31)\\
&= \lim_{n\to\infty} \operatorname{Tr}\left(e^{t_1 L_{s*}}(\tilde{\rho})\rho_{ss}^{1/4} e^{t_2\mathcal{L}_s}\left(P_n B P_n\right)\rho_{ss}^{1/4}\right). && (32)
\end{aligned}$$

Similar to before, we define $B_n = P_n B P_n$ where $B = e^{(t_3+\ldots+t_l)\mathcal{L}_s}(\mathbf{1})$. By repeating the above argument until all the t_i have been exhausted we arrive at the desired expression

$$\mathrm{Tr}\left(e^{(t_1+\ldots+t_l)L_{s*}}(\tilde{\rho})\rho_{ss}^{1/2}\right) = \mathrm{Tr}\left(\tilde{\rho}e^{tL_s}(\rho_{ss}^{1/2})\right).$$

This shows that the semigroup L_s on the Hilbert-Schmidt space gives rise to the desired expectation values.